

## Modeling impacts of farming management alternatives on CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O emissions: A case study for water management of rice agriculture of China

Changsheng Li,<sup>1</sup> Steve Frolking,<sup>1</sup> Xiangming Xiao,<sup>1</sup> Berrien Moore III,<sup>1</sup> Steve Boles,<sup>1</sup> Jianjun Qiu,<sup>2</sup> Yao Huang,<sup>3</sup> William Salas,<sup>4</sup> and Ronald Sass<sup>5</sup>

Received 16 July 2004; revised 10 March 2005; accepted 9 May 2005; published 17 August 2005.

[1] Since the early 1980s, water management of rice paddies in China has changed substantially, with midseason drainage gradually replacing continuous flooding. This has provided an opportunity to estimate how a management alternative impacts greenhouse gas emissions at a large regional scale. We integrated a process-based model, DNDC, with a GIS database of paddy area, soil properties, and management factors. We simulated soil carbon sequestration (or net CO<sub>2</sub> emission) and CH<sub>4</sub> and N<sub>2</sub>O emissions from China's rice paddies (30 million ha), based on 1990 climate and management conditions, with two water management scenarios: continuous flooding and midseason drainage. The results indicated that this change in water management has reduced aggregate CH<sub>4</sub> emissions about 40%, or 5 Tg CH<sub>4</sub> yr<sup>-1</sup>, roughly 5–10% of total global methane emissions from rice paddies. The mitigating effect of midseason drainage on CH<sub>4</sub> flux was highly uneven across the country; the highest flux reductions (>200 kg CH<sub>4</sub>-C ha<sup>-1</sup> yr<sup>-1</sup>) were in Hainan, Sichuan, Hubei, and Guangdong provinces, with warmer weather and multiple-cropping rice systems. The smallest flux reductions (<25 kg CH<sub>4</sub>-C ha<sup>-1</sup> yr<sup>-1</sup>) occurred in Tianjin, Hebei, Ningxia, Liaoning, and Gansu Provinces, with relatively cool weather and single cropping systems. Shifting water management from continuous flooding to midseason drainage increased N<sub>2</sub>O emissions from Chinese rice paddies by 0.15 Tg N yr<sup>-1</sup> (~50% increase). This offset a large fraction of the greenhouse gas radiative forcing benefit gained by the decrease in CH<sub>4</sub> emissions. Midseason drainage-induced N<sub>2</sub>O fluxes were high (>8.0 kg N/ha) in Jilin, Liaoning, Heilongjiang, and Xinjiang provinces, where the paddy soils contained relatively high organic matter. Shifting water management from continuous flooding to midseason drainage reduced total net CO<sub>2</sub> emissions by 0.65 Tg CO<sub>2</sub>-C yr<sup>-1</sup>, which made a relatively small contribution to the net climate impact due to the low radiative potential of CO<sub>2</sub>. The change in water management had very different effects on net greenhouse gas mitigation when implemented across climatic zones, soil types, or cropping systems. Maximum CH<sub>4</sub> reductions and minimum N<sub>2</sub>O increases were obtained when the mid-season draining was applied to rice paddies with warm weather, high soil clay content, and low soil organic matter content, for example, Sichuan, Hubei, Hunan, Guangdong, Guangxi, Anhui, and Jiangsu provinces, which have 60% of China's rice paddies and produce 65% of China's rice harvest.

**Citation:** Li, C., S. Frolking, X. Xiao, B. Moore III, S. Boles, J. Qiu, Y. Huang, W. Salas, and R. Sass (2005), Modeling impacts of farming management alternatives on CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O emissions: A case study for water management of rice agriculture of China, *Global Biogeochem. Cycles*, 19, GB3010, doi:10.1029/2004GB002341.

<sup>1</sup>Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, New Hampshire, USA.

<sup>2</sup>Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Science, Beijing, China.

<sup>3</sup>College of Resource and Environmental Sciences, Nanjing Agricultural University, Nanjing, China.

<sup>4</sup>Applied Geosolutions, LLC, Durham, New Hampshire, USA.

<sup>5</sup>Department of Ecology and Evolutionary Biology, Rice University, Houston, Texas, USA.

### 1. Introduction

[2] Food production contributes approximately 70% of global atmospheric input of nitrous oxide (N<sub>2</sub>O) and 40% of global atmospheric input of methane (CH<sub>4</sub>) [Cole *et al.*, 1996], and so represents a significant opportunity for greenhouse gas mitigation through reductions of CH<sub>4</sub> and N<sub>2</sub>O emissions, as well as through soil carbon sequestration [Oenema *et al.*, 2001]. When assessing the impact of food and fiber production systems on the Earth's radiation budget, the entire suite of greenhouse gases (i.e., CO<sub>2</sub>,

CH<sub>4</sub>, and N<sub>2</sub>O) needs to be considered [Li, 1995; Robertson *et al.*, 2000; Smith *et al.*, 2001; Li *et al.*, 2005]. Since each greenhouse gas has its own radiative potential [Ramaswamy *et al.*, 2001], a net global warming potential (GWP) of a crop production system can be estimated, accounting for all the three gases.

[3] Rice is a major food crop in Asia (~130 million hectares were sown in 2002 [Food and Agriculture Organization (FAO), 2004]), and the majority of rice production in Asia is from flooded paddy fields (<10% of sown area is upland rice [Huke and Huke, 1997]). Rice paddies contribute about 10% of total global methane emissions to the atmosphere [Prather *et al.*, 2001]. Field studies have shown that water management can have a significant influence on total methane emissions during a cropping season [Wassmann *et al.*, 2000a; Sass *et al.*, 1992] so paddy water management has become a target mitigation scenario [Wassmann *et al.*, 2000b].

[4] Water management in paddy fields can be classified as irrigated or rainfed [Huke and Huke, 1997]. Irrigated paddies typically have continuous management of their flooding regime, while rainfed paddy flooding can be sporadic, depending on seasonal precipitation. Water management of irrigated rice paddies can be coarsely partitioned into two categories: continuous flooding or mid-season draining. In continuous flooding, the paddy soils remain saturated and puddled from just prior to transplanting until just before harvest. Puddled paddy soils quickly become anoxic, and remain so for the duration of the growing season. Mid-season draining/drying entails active draining of the paddy or passive drying for 1 to 2 weeks, typically several times during a growing season. During this period the surface soil becomes more oxidic. Field studies have shown that mid-season draining reduces total crop-season CH<sub>4</sub> emissions by 10–80% [Sass *et al.*, 1992; Yagi *et al.*, 1996; Cai *et al.*, 1999; Wassmann *et al.*, 2000a], giving the practice a strong potential for greenhouse gas mitigation [Wassmann *et al.*, 2000b]. Fewer field studies have measured the consequences of mid-season draining on N<sub>2</sub>O emissions but there are strong indications that mid-season draining can cause an increase in N<sub>2</sub>O flux [e.g., Chen *et al.*, 1995; Zheng *et al.*, 1997, 2000], probably because the oxidic/anoxic transitions favor both nitrification and denitrification.

[5] Mid-season drainage was initiated in rice farming in northeastern China in the early 1980s owing to frequent shortages of irrigation water. Mid-season drainage was found to not only reduce water use but also increase crop yield. The practice was quickly adopted for rice agriculture in northern China in the 1980s, and spread through most of China's rice agriculture during the 1990s [Shen *et al.*, 1998]. Over the years, mid-season drainage has demonstrated other agronomic advantages, including reducing ineffective tillers, removing toxic substances, and maintaining healthier roots under anaerobic soil conditions. Short periods of drainage for soil aeration during vegetative growth and intermittent irrigation during reproductive growth are now common in China [Gao *et al.*, 1992] and Japan [Yoshida, 1981]. The intensity of drainage and the interval between the cycles of flooding-

drainage-reflooding vary with soil characteristics and weather conditions.

[6] Water management for rice agriculture in China, most of which is irrigated [Huke and Huke, 1997], has changed substantially during the 20-year period from 1980 to 2000, with midseason drainage gradually replacing continuous flooding [Shen *et al.*, 1998]. This nationwide change provided an opportunity for us to quantify how a management alternative could impact greenhouse gas emissions at a large regional scale. In an earlier study we evaluated the impact of this changing management on methane emissions from China's rice paddies, and estimated a reduction of about 40% in total emissions from all paddy rice in China [Li *et al.*, 2002]. In this paper we extend this analysis to look at changes in both CH<sub>4</sub> and N<sub>2</sub>O emissions, and to look at spatial patterns of changing greenhouse gas across China, rather than just an aggregate national total.

## 2. Methods

[7] Both CH<sub>4</sub> and N<sub>2</sub>O fluxes from agro-ecosystems are highly variable in space and time, affected by ecological drivers (e.g., climate, vegetation, and anthropogenic activity), soil environmental factors (e.g., temperature, moisture, pH, redox potential, and substrate concentration gradients), and biochemical or geochemical reactions [Li, 2000, 2001; Li *et al.*, 2004]. Process-based models are used to quantify trace gas fluxes driven by the local climate, soil, vegetation, and management conditions at the site scale. GIS databases provide spatially differentiated information of climate, soil, vegetation, and management to drive the model runs across the region. To quantify the impacts of the water management change on C sequestration and CH<sub>4</sub> and N<sub>2</sub>O emissions from ~30 million hectares of rice paddies in China, we integrated a process-based model, DNDC, with a GIS database of paddy area, soil properties, paddy management (fertilizer use, water management, crop residue management, planting and harvest dates), and daily weather data.

### 2.1. DNDC Model

[8] DNDC was originally developed for predicting carbon sequestration and trace gas emissions for non-flooded agricultural lands, simulating the fundamental processes controlling the interactions among ecological drivers, soil environmental factors, and relevant biochemical or geochemical reactions, which collectively determine the rates of trace gas production and consumption in agricultural ecosystems [Li *et al.*, 1992, 1994, 1996]. Details of management (e.g., crop rotation, tillage, fertilization, manure amendment, irrigation, weeding, and grazing) have been parameterized and linked to the various biogeochemical processes (e.g., crop growth, litter production, soil water infiltration, decomposition, nitrification, denitrification) embedded in DNDC. To enable DNDC to simulate C and N biogeochemical cycling in paddy rice ecosystems, we modified the model by adding a series of anaerobic processes. The paddy-rice version of DNDC has been described and tested in recent manuscripts [Li *et al.*, 2002; Cai *et al.*, 2003; Li *et al.*, 2004], and is summarized briefly here.

**Table 1.** DNDC Oxidation-Reduction Scheme


Dominant Oxidant	Eh Range, mV	Dominant Reactions Simulated
O <sub>2</sub>	650–500	aerobic decomposition, CH <sub>4</sub> oxidation, nitrification
NO <sub>3</sub> <sup>-</sup> , NO <sub>2</sub> <sup>-</sup> , NO, N <sub>2</sub> O	500–200	denitrification sequence
Mn <sup>3+</sup>	200–100	manganese reduction
Fe <sup>3+</sup>	100–0	iron reduction
SO <sub>4</sub> <sup>2-</sup>	0––150	sulfate reduction
H <sub>2</sub>	–150––350	methanogenesis

[9] Paddy soil is characterized by the frequent changes between saturated and unsaturated conditions driven by water management. During these changes in soil water content, the soil redox potential (i.e., Eh) is subject to substantial changes between +600 and –300 mV. One of the key processes controlling CH<sub>4</sub> and N<sub>2</sub>O production/consumption in paddy soils is soil Eh dynamics; CH<sub>4</sub> or N<sub>2</sub>O are produced or consumed under certain Eh conditions (–300 to –150 mV for CH<sub>4</sub>, and 200 to 500 mV for N<sub>2</sub>O), so the two gases are produced during different stages of the varying soil redox potential.

[10] To quantify Eh dynamics and its impacts on N<sub>2</sub>O and CH<sub>4</sub> production, DNDC combines the Nernst equation, a basic thermodynamic formula defining soil Eh based on concentrations of the existing oxidants and reductants in the soil liquid phase [Stumm and Morgan, 1981], with the Michaelis-Menten equation, a widely applied formula describing kinetics of microbial growth with dual nutrients [Paul and Clark, 1989]. The Nernst and the Michaelis-Menten equations can be linked in model calculations since the two equations share a common factor, oxidant concentration. A simple kinetic scheme was adopted in DNDC to define the effective anaerobic volumetric fraction of a soil, based on the Eh value as calculated with the Nernst equation, using the concentration of the dominant active oxidant (in order of descending Eh: O<sub>2</sub>, NO<sub>3</sub><sup>-</sup>, Mn<sup>4+</sup>, Fe<sup>3+</sup>, SO<sub>4</sub><sup>2-</sup>, CO<sub>2</sub>). Each oxidant is assigned an Eh range (Table 1). When a soil layer's Eh value is in a particular oxidant's range, the layer is divided into a high Eh zone (Eh = range upper value) and a low Eh zone (Eh = range lower value), as a linear function of where within the range the Eh value is. DNDC allocates substrates (e.g., DOC, NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup>, CH<sub>4</sub>) to reductive reactions (e.g., denitrification, methanogenesis) and oxidative reactions (e.g., nitrification, methanotrophy) based on relative fractional volumes of the oxidizing and reducing zones, and the potential oxidation and reduction reactions are determined by Eh (and pH) [Stumm and Morgan, 1981]. When a soil is flooded, oxygen diffusion into the soil is severely restricted, and ongoing decomposition lowers the oxygen concentration, reducing the soil Eh and causing the low Eh volume fraction to increase. As the oxygen is depleted, the low Eh volume fraction reaches its maximum. At this moment, a new oxidant (i.e., NO<sub>3</sub><sup>-</sup>) will become the dominant species in the soil, and a new, lower Eh volume fraction will begin to swell, driven by NO<sub>3</sub><sup>-</sup> depletion. By tracking the formation and deflation of a series of Eh volume fractions driven by depletions of O<sub>2</sub>, NO<sub>3</sub><sup>-</sup>, Mn<sup>4+</sup>, Fe<sup>3+</sup>, and SO<sub>4</sub><sup>2-</sup>, consecutively, DNDC estimates soil Eh dynamics as well as rates of reductive/


oxidative reactions, which produce and consume CH<sub>4</sub> or N<sub>2</sub>O in the soil. This links the soil water regime to trace gas emissions for rice paddy ecosystems, and DNDC predicts daily CH<sub>4</sub> and N<sub>2</sub>O fluxes from rice fields through the growing and fallow seasons, as they remain flooded or shift between flooded and drained.

[11] This new DNDC model has been tested against several methane flux data sets from wetland rice sites in the United States, Italy, China, Thailand, and Japan [Li *et al.*, 2002; Cai *et al.*, 2003]. For sites in East Asia, simulated seasonal CH<sub>4</sub> emissions from paddy soils were in good agreement with field studies ( $r^2 = 0.96$ , regression slope = 1.1,  $n = 23$ , range in fluxes 9 to 725 kg CH<sub>4</sub>-C ha<sup>-1</sup> season<sup>-1</sup>) [Cai *et al.*, 2003]. There were fewer sites with N<sub>2</sub>O flux data. If the site soil characteristics and crop and water management were well described, simulated fluxes were similar to observation (Figure 1) [Zheng *et al.*, 1997]. However, for the three cases where the management details were not so well known, there were discrepancies between model and field, though there was overlap in the range in fluxes, 0.6 to 2.0 kg N<sub>2</sub>O-N ha<sup>-1</sup> season<sup>-1</sup> for the field and 0.4 to 5.7 kg N<sub>2</sub>O-N ha<sup>-1</sup> season<sup>-1</sup> for the model [Cai *et al.*, 2003]. The results from the tests indicate that DNDC is capable of estimating the seasonal magnitudes of CH<sub>4</sub> and N<sub>2</sub>O fluxes from paddy sites, although discrepancies exist for about 20% of the tested cases.



**Figure 1.** Comparison between observed and DNDC-modeled CH<sub>4</sub> and N<sub>2</sub>O fluxes from a paddy rice field applied with mid-season drainage in Wu County, Jiangsu Province, China, in 1995. DNDC captured the episodes of CH<sub>4</sub> emission depressions and N<sub>2</sub>O emission increases during the soil drying time periods by tracking the soil oxygen diffusion, CH<sub>4</sub> oxidation, labile organic matter decomposition, and stimulated nitrification and denitrification fueled by the increased ammonium and nitrate production due to the conversions of soil anaerobic to aerobic conditions driven by the mid-season drainage. (Field data were adopted from Zheng *et al.* [1997]). See color version of this figure at back of this issue.





**Figure 2.** Mean annual provincial rough rice production for 1991–2000 (IRRI 2004) in millions of tonnes per year. No provincial data were reported for Tibet (Xizang), Qinghai, Xinjiang, Gansu, Shanxi, Inner Mongolia, and Beijing; 1991–2000 total mean annual rough rice production for these seven provinces was  $740,000 \text{ t yr}^{-1}$ . Taiwan was excluded from our entire analysis because of a lack of crop rotation data.

## 2.2. GIS Database

[12] The model domain contained 30 million ha of rice paddies in China [Frolking *et al.*, 2002]. There were 11 different crop rotations, including single-rice, double-rice, rice-winter wheat, rice-rapeseed, rice-rice-vegetable, etc. The area occupied by each rotation in each county was quantified by combining a county-scale statistical database of crop-sown areas with a Landsat TM-derived land-cover map for all of mainland China [Frolking *et al.*, 2002]. The majority of rice production occurs in southern China, and particularly along the Yangtze River (Figure 2 [International Rice Research Institute (IRRI), 2004]). Daily weather data (maximum and minimum air temperatures, precipitation) for 1990 from 610 weather stations across China were acquired from the National Center for Atmospheric Research (<http://dss.ucar.edu/index.html>). Station data were assigned to each county on a nearest neighbor basis. Maximum and minimum values of soil texture, pH, bulk density, and organic carbon content were derived for each county from digitization of national soil maps [Institute of Soil Science, 1986] and other information [National Soil Survey Office of China, 1997]. Model simulations could then be conducted for each county by choosing soil parameter values spanning the observed range. Soil Mn, Fe, and sulfate contents were set to average values for Chinese paddy soils: Mn =  $30 \text{ mg kg}^{-1}$  soil, Fe =  $80 \text{ mg kg}^{-1}$  soil, and sulfate =  $220 \text{ mg kg}^{-1}$  soil [Li, 1992]. General data on fertilizer use, tillage, planting and harvest dates, crop residue management, and crop varieties were taken from Central Radio and Television School of Agriculture (CRTSA) [1995], Huang *et al.*

[1997], Cui *et al.* [1994], Liu and Mu [1993], and Beijing Agricultural University [1992]. Shen [1998] reported that based on national statistics, an average of 30% of total crop residue (leaves + stems + roots) was returned to the soil, which we adopted for all fields. Manure production was based on animal and human populations from the county database assembled by the Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, using standard manure production rates [Intergovernmental Panel on Climate Change (IPCC), 1997], and field application rates of 20% for animal manure and 10% for human manure.

[13] Changing water management of China's rice paddies is not well documented. Li *et al.* [2002] constructed a simple scenario of the evolution of paddy water management in China from 2000, as paddies went from near 100% continuous flooding in 1980 [Shen *et al.*, 1998] to ~80% midseason drainage by 2000 (Qingmu Chen, Chinese Academy of Agricultural Sciences, personal communication, 2002). In this study we wish to quantify the potential impacts of alternative water management practices, so we designed two water management scenarios: continuous flooding (CF) and midseason drainage (MSD) for 100% of the rice paddies in China.

## 2.3. Uncertainty

[14] The county was chosen as the basic spatial unit for GIS database construction since most of the statistical cropland data was county-based. Meteorological data, soil properties, and agricultural management data were obtained from ground-based sources. Since each county is regarded to be uniform during a single model simulation, uncertainty estimates related to the inherent heterogeneities of many input parameters within the county must be generated during the scaling-up process. With ~2500 counties, most with rice crops and many with several crop rotations, a set of annual simulations with a single parameter set required about 10,000 model runs, so Monte Carlo analysis (randomly adjusting all parameters thousands of times to get a statistical distribution of outputs) was computationally prohibitive.

[15] Instead, sensitivity tests were conducted to prioritize the environmental factors regarding their effects on  $\text{CH}_4$  or  $\text{N}_2\text{O}$  emissions [Li *et al.*, 2004]. Among the tested factors, including soil properties, temperature, and precipitation, the most sensitive factors for  $\text{CH}_4$  and  $\text{N}_2\text{O}$  emissions were soil texture and soil organic carbon (SOC) content, respectively. Varying soil texture and SOC over the ranges reported in the county-scale database produced a range of  $\text{CH}_4$  and  $\text{N}_2\text{O}$  emissions. The range in fluxes generated by this “most sensitive factor” method was compared to the distribution of fluxes generated by a Monte Carlo simulation for three paddy sites (Yunnan Province, China; Suphan Buri Province, Thailand; and California, United States); 70–98% of annual  $\text{CH}_4$  fluxes and 61–88% of annual  $\text{N}_2\text{O}$  fluxes generated in 5000 Monte Carlo simulations for each site fell within the range generated by varying the most sensitive factor [Li *et al.*, 2004].

[16] Applying the “most sensitive factor” method, we ran DNDC for each rice rotation in each county twice, once

**Table 2.** Provincial Paddy Area and Simulated CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O Maximum and Minimum Fluxes for Continuous Flooding (CF) and Mid-Season Draining (MSD) Water Management<sup>a</sup>

Province <sup>b</sup>	Production, <sup>c</sup> Gg C yr <sup>-1</sup>	CO <sub>2</sub> , Gg C yr <sup>-1</sup>				CH <sub>4</sub> , Gg C yr <sup>-1</sup>				N <sub>2</sub> O, Gg N yr <sup>-1</sup>			
		CF <sub>min</sub>	CF <sub>max</sub>	MSD <sub>min</sub>	MSD <sub>max</sub>	CF <sub>min</sub>	CF <sub>max</sub>	MSD <sub>min</sub>	MSD <sub>max</sub>	CF <sub>min</sub>	CF <sub>max</sub>	MSD <sub>min</sub>	MSD <sub>max</sub>
Sichuan <sup>d</sup>	11,000	1500	240	1600	-380	1800	1300	1300	200	49	34	72	47
Hunan	12,000	370	-1700	280	-2000	1100	550	580	110	38	32	54	41
Jiangsu	9000	1800	1400	1900	1300	1200	770	1000	480	25	18	36	25
Jiangxi	7600	74	-1000	-3	-1200	1100	560	710	72	33	22	54	31
Anhui	6300	600	-710	780	-920	1100	460	870	77	29	14	38	19
Hubei	8300	510	-1200	490	-1300	1100	410	650	46	24	13	33	16
Guangdong	7600	310	-700	57	-870	860	470	400	150	26	22	40	29
Guangxi	6200	1800	15	1700	-13	670	260	260	13	38	26	56	33
Yunnan	2600	180	-290	130	-320	160	81	18	3	12	12	17	14
Guizhou	2200	450	-560	530	-550	200	96	74	9	14	10	21	13
Fujian	3600	-1	-180	-39	-250	260	170	140	29	14	10	21	13
Henan	1600	340	88	300	26	340	140	230	27	6	3	9	4
Liaoning	1800	490	390	660	450	1800	140	230	61	10	8	19	17
Jilin	1700	650	370	680	500	150	100	120	30	12	8	22	24
Shaanxi	450	130	180	120	160	110	64	77	7	4	3	7	4
Shanghai	800	160	140	160	120	150	95	120	56	2	2	3	2
Inner Mongolia	n.a. <sup>e</sup>	120	160	120	180	34	21	27	5	3	2	4	3
Shandong	520	-47	20	-44	27	61	50	50	34	1	1	2	2
Hebei	450	-22	22	-5	37	28	26	34	27	1	1	1	1
Zhejiang	6100	340	-390	340	-480	730	390	550	140	23	17	34	23
Heilongjiang	3200	1800	1400	1800	1600	270	180	230	78	36	26	44	44
Tianjin	190	-14	11	7	27	17	16	32	26	0.7	0.6	1	1
Ningxia	270	-47	-38	-29	-36	2	3	10	2	0.7	0.5	1	1
Gansu	n.a. <sup>e</sup>	7	11	7	12	2	2	2	0.6	0.1	0.1	0.2	0.2
Hainan	790	610	210	610	200	200	100	73	19	8	4	12	5
Beijing	n.a. <sup>e</sup>	-1	7	3	7	5	4	6	2	0.1	0.1	0.2	0.1
Shanxi	n.a. <sup>e</sup>	4	7	6	8	3	3	3	2	0.1	0.1	0.2	0.2
Xinjiang	n.a. <sup>e</sup>	51	56	57	63	51	27	42	7	2	1	2	2
<b>China</b>	<b>94,000</b>	<b>12,000</b>	<b>-2000</b>	<b>12,000</b>	<b>-3500</b>	<b>12,000</b>	<b>6400</b>	<b>7800</b>	<b>1700</b>	<b>410</b>	<b>290</b>	<b>610</b>	<b>420</b>

<sup>a</sup>CO<sub>2</sub> flux equals negative of change in soil organic carbon storage. Sign convention for all fluxes is positive flux equals net emission from soil. All areas and all flux values greater than 10 Gg were rounded to two significant figures.

<sup>b</sup>Provinces are sorted by paddy area (see Table 3). Tibet and Qinghai Provinces are excluded due to small paddy areas.

<sup>c</sup>Mean annual rough rice production is given for 1991–2000 [IRRI, 2004]; carbon fraction of dry weight is assumed to be 50%.

<sup>d</sup>This includes Chongqing Province, which was established from land in Sichuan Province in 1997.

<sup>e</sup>Production data (see note 2 above) are not reported individually for some provinces; total production for all non-reported provinces = 370 Gg C yr<sup>-1</sup>; China total equals sum of all reported and non-reported provinces.

with low SOC, low pH, and high clay content, and once with high SOC, high pH, and low clay content, to produce a range in CH<sub>4</sub> and N<sub>2</sub>O emissions wide enough to represent likely variations in actual fluxes caused by the heterogeneous nature of soil properties.

#### 2.4. Emissions Calculations

[17] Of 11 possible paddy rotations mapped across China [Frolking *et al.*, 2002], a single county typically had 2–4 paddy rotations. For each paddy crop rotation in each county, we simulated annual change in SOC ( $\Delta$ SOC), and annual CH<sub>4</sub> and N<sub>2</sub>O emissions for the range of soil conditions (as discussed in section 2.3). We consider  $\Delta$ SOC as the negative of the site's net annual CO<sub>2</sub> flux. Methane flux is another pathway for carbon out of wetland soil, but the substrates for methane production are primarily plant-derived C (e.g., root exudation, deposition, and respiration CO<sub>2</sub>) [Watanabe *et al.*, 1999; Lu *et al.*, 2002; King *et al.*, 2002], which are not included in the simulation's SOC budget, so methane flux would only be a small fraction of simulated  $\Delta$ SOC. We then summed the low and high emissions from all of the rice cropping systems in the county, based on each

rotation's fraction of total cropland area, to get a range in total CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O emissions for the county. We calculated a change in flux due to changing water management from continuous flooding to mid-season draining as the difference between the mean MSD flux and the mean CF flux, where the mean flux is the average of the high and low fluxes.

[18] The climate impact of these flux changes can be compared by converting them to a common basis of CO<sub>2</sub>-equivalent fluxes using the global warming potential (GWP) methodology [Ramaswamy *et al.*, 2001]. GWP values for CH<sub>4</sub> and N<sub>2</sub>O depend on the time horizon chosen. Methane, with a much shorter atmospheric adjustment time than N<sub>2</sub>O or CO<sub>2</sub>, has very different 20-year, 100-year, and 500-year GWP values [Ramaswamy *et al.*, 2001]. A CO<sub>2</sub>-equivalent flux for a 1-kg CH<sub>4</sub> or N<sub>2</sub>O flux is the mass of CO<sub>2</sub> emissions that would generate the same integrated radiative forcing, over the chosen time horizon, as a 1-kg emission CH<sub>4</sub> or N<sub>2</sub>O. (Note that GWP calculations are done with molecular masses, not C or N masses.) For example, let  $\Delta$ CO<sub>2</sub>,  $\Delta$ CH<sub>4</sub>, and  $\Delta$ N<sub>2</sub>O be the differences between the mean carbon dioxide, methane, and nitrous oxide fluxes (MSD minus CF) in kg C or N ha<sup>-1</sup> yr<sup>-1</sup>. Then the 20-year

time horizon CO<sub>2</sub>-equivalent flux, in kg CO<sub>2</sub> ha<sup>-1</sup> yr<sup>-1</sup>, is calculated as


$$\begin{aligned} (\text{CO}_{2\text{-eq.}})_{20} &= \Delta[\text{CO}_2 \text{ flux}] + \Delta[\text{methane's CO}_2 \text{ - equivalent}] \\ &\quad + \Delta[\text{nitrous oxide's CO}_2 \text{ - equivalent}] \\ &= \Delta\text{CO}_2 \cdot \frac{44}{12} + \Delta\text{CH}_4 \cdot \frac{16}{12} \cdot \text{GWP}_{20}^{\text{CH}_4} + \Delta\text{N}_2\text{O} \cdot \frac{44}{28} \\ &\quad \cdot \text{GWP}_{20}^{\text{N}_2\text{O}}, \end{aligned} \quad (1)$$

where the numerical fractions in the second line are the ratios of molecular weights (CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O) to elemental weights (C and N); GWP<sub>20</sub><sup>CH<sub>4</sub></sup> equals 62 kg CO<sub>2</sub>-eq kg<sup>-1</sup> CH<sub>4</sub> and GWP<sub>20</sub><sup>N<sub>2</sub>O</sup> equals 275 kg CO<sub>2</sub>-eq kg<sup>-1</sup> N<sub>2</sub>O; GWP<sub>100</sub><sup>CH<sub>4</sub></sup> equals 23 kg CO<sub>2</sub>-eq kg<sup>-1</sup> CH<sub>4</sub> and GWP<sub>100</sub><sup>N<sub>2</sub>O</sup> equals 296 kg CO<sub>2</sub>-eq kg<sup>-1</sup> N<sub>2</sub>O; GWP<sub>500</sub><sup>CH<sub>4</sub></sup> equals 7 kg CO<sub>2</sub>-eq kg<sup>-1</sup> CH<sub>4</sub>, and GWP<sub>500</sub><sup>N<sub>2</sub>O</sup> equals 156 kg CO<sub>2</sub>-eq kg<sup>-1</sup> N<sub>2</sub>O [Ramaswamy *et al.*, 2001].

### 3. Results

[19] The ranges in total CO<sub>2</sub> emissions with CF and MSD managements were +12,000 to -2000 and +12,000 to -3500 Gg CO<sub>2</sub>-C yr<sup>-1</sup>, respectively (Table 2). A positive CO<sub>2</sub> emission corresponds to a net loss of soil organic carbon, so overall mean behavior was probably an SOC loss in both scenarios though zero net flux fell within both predicted ranges. The CO<sub>2</sub> flux range went more negative (i.e., CO<sub>2</sub> uptake) for the MSD scenario, implying a small shift toward lower CO<sub>2</sub> emissions. The shift on CO<sub>2</sub> emission between the two water management scenarios was caused by changes in soil decomposition rates and crop residue incorporation rates. The latter was affected by soil N availability and total crop biomass production, which were both affected by water management. Taking the mean values, changing water management from continuous flooding to midseason draining caused a net uptake (i.e., reduced loss rate) of CO<sub>2</sub> of 640 Gg CO<sub>2</sub>-C yr<sup>-1</sup>. For perspective, in 2000, fossil fuel carbon emissions from China were about 700,000 Gg C yr<sup>-1</sup> [Marland and Boden, 2004]. Changing water management to MSD led overall to net losses of CO<sub>2</sub> in northern China, and net uptake of CO<sub>2</sub> in southern China (Figure 3a), but changes were generally not large (Table 2).

[20] The ranges in total CH<sub>4</sub> emissions with CF and MSD managements were 6400–12,000 and 1700–7800 Gg CH<sub>4</sub>-C yr<sup>-1</sup>, respectively (Table 2). Taking the mean values, changing water management from continuous flooding to midseason draining caused a reduction in aggregate methane emissions from rice paddies of about 40%, or 6000 Gg CH<sub>4</sub> yr<sup>-1</sup>. This is 5–10% of total global rice paddy methane emissions [Prather *et al.*, 2001]. The mitigating effect of MSD on CH<sub>4</sub> flux was highly uneven across the country.



**Figure 3.** Change, due to conversion from continuous flooding to mid-season draining, in annual paddy flux rates of (a) CO<sub>2</sub>, (b) CH<sub>4</sub>, and (c) N<sub>2</sub>O. Each provincial value is the difference between the mean for all counties of high and low values for mid-season draining and the mean for all counties of high and low values for continuous flooding. Sign convention: a positive flux difference (red shades) represents an increase in flux to the atmosphere, and a negative flux difference (blue shades) represents a decrease in flux to the atmosphere. For all provinces, the water management conversion reduced mean CH<sub>4</sub> flux and increased mean N<sub>2</sub>O flux. Tibet (Xizang) and Qinghai Provinces have very small paddy areas, and are reported here as no data; Taiwan was excluded from the analysis because of a lack of crop rotation data. See color version of this figure at back of this issue.



**Table 3.** Provincial Rice Production and CO<sub>2</sub>-Equivalent Fluxes for Greenhouse Gas Impact of Changing Management From Continuous Flooding to Mid-Season Drainage (MSD Mean Flux Minus CF Mean Flux), Using 20-, 100-, and 500-Year Time Horizons<sup>a</sup>

Province <sup>b</sup>	Paddy Area, <sup>c</sup> ha	20-Year CO <sub>2</sub> Equivalent, kg CO <sub>2</sub> ha <sup>-1</sup> yr <sup>-1</sup>				100-Year CO <sub>2</sub> Equivalent, kg CO <sub>2</sub> ha <sup>-1</sup> yr <sup>-1</sup>				500-Year CO <sub>2</sub> Equivalent, kg CO <sub>2</sub> ha <sup>-1</sup> yr <sup>-1</sup>			
		Total	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	Total	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	Total	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
Sichuan <sup>d</sup>	3,700,000	-16,000	270	-18,000	2100	-4200	270	-6700	2200	-610	270	-2100	1200
Hunan	3,200,000	-11,000	180	-13,000	1700	-2800	180	-4800	1800	-330	180	-1500	940
Jiangsu	2,500,000	-6000	-5	-7600	1600	-1100	-5	-2800	1700	+22	-5	-850	880
Jiangxi	2,500,000	-11,000	210	-14,000	2600	-2300	210	-5300	2800	+70	210	-1600	1500
Anhui	2,000,000	-12,000	19	-13,000	1400	-3300	19	-4800	1500	-630	19	-1500	820
Hubei	1,900,000	-17,000	110	-18,000	1400	-5200	110	-6700	1500	-1200	110	-2100	770
Guangdong	1,900,000	-14,000	410	-17,000	2400	-3300	410	-6300	2500	-150	410	-1900	1300
Guangxi	1,700,000	-12,000	83	-16,000	3100	-2400	83	-5800	3400	+95	83	-1800	1800
Yunnan	1,500,000	-4600	94	-5900	1200	-840	94	-2200	1200	+85	94	-660	650
Zhejiang	1,200,000	-9100	100	-12,000	2500	-1500	100	-4400	2700	+220	100	-1300	1400
Heilongjiang	1,000,000	-460	-360	-4300	4200	+2500	-360	-1600	4500	+1500	-360	-480	2400
Guizhou	710,000	-5500	-130	-7200	1800	-830	-130	-2700	2000	+98	-130	-820	1000
Fujian	690,000	-8700	210	-11,000	2200	-1600	210	-4100	2400	+200	210	-1300	1200
Henan	650,000	-12,000	270	-13,000	1200	-3300	270	-4900	1300	-530	270	-1500	690
Liaoning	280,000	+3300	-630	-1800	5600	+4800	-630	-650	6100	+2400	-630	-200	3200
Jilin	240,000	+1800	-470	-6200	8400	+6300	-470	-2300	9100	+3600	-470	-700	4800
Hainan	190,000	-20,000	55	-24,000	3200	-5300	55	-8800	3500	-800	55	-2700	1800
Shaanxi	180,000	-11,000	140	-14,000	3300	-1500	140	-5200	3500	+400	140	-1600	1800
Shanghai	150,000	-9600	110	-11,000	1500	-2500	110	-4200	1600	-330	110	-1300	830
Inner Mongolia	150,000	-1900	-170	-4800	3000	+1300	-170	-1800	3300	+1000	-170	-540	1700
Shandong	140,000	-3700	-110	-6000	2400	+270	-110	-2200	2600	+590	-110	-680	1400
Hebei	100,000	+2700	-390	1800	1300	+1700	-390	680	1400	+540	-390	210	730
Xinjiang	93,000	-6300	-200	-10,000	4100	+460	-200	-3800	4500	+990	-200	-1200	2300
Tianjin	38,000	+10,000	-660	9800	840	+3900	-660	3600	900	+920	-660	1100	470
Ningxia	38,000	+3800	-390	3000	1200	+2000	-390	1100	1300	+610	-390	340	660
Gansu	23,000	-870	-44	-2000	1100	+460	-44	-730	1200	+380	-44	-220	650
Beijing	16,000	-460	-320	-1100	960	+310	-320	-410	1000	+100	-320	-120	540
Shanxi	11,000	-1700	-280	-4100	2700	+1100	-280	-1500	2900	+770	-280	-460	1500
<b>China<sup>e</sup></b>	<b>30,000,000</b>	<b>-9900</b>	<b>79</b>	<b>-12,000</b>	<b>2300</b>	<b>-2000</b>	<b>79</b>	<b>-4600</b>	<b>2500</b>	<b>+6</b>	<b>79</b>	<b>-1400</b>	<b>1300</b>

<sup>a</sup>The CO<sub>2</sub>-equivalent flux was calculated as in equation (1) in the text. A positive CO<sub>2</sub>-eq. flux means a net emission to the atmosphere, and a negative flux means a net uptake from the atmosphere. All production values and all flux values greater than 10 kg CO<sub>2</sub>-eq. ha<sup>-1</sup> yr<sup>-1</sup> were rounded to two significant figures.

<sup>b</sup>Tibet and Qinghai Provinces (not included) have little paddy rice, and generate <1 Gg CO<sub>2</sub>-C yr<sup>-1</sup>, <0.1 Gg CH<sub>4</sub>-C yr<sup>-1</sup>, and <0.1 Gg N<sub>2</sub>O-N yr<sup>-1</sup>.

<sup>c</sup>This comprises paddy land area, and does not double-count double rice crops; from *Frolking et al.* [2002].

<sup>d</sup>This includes Chongqing Province, which was established from land in Sichuan Province in 1997.

<sup>e</sup>China totals equal sum of all reported and nonreported provinces.

The highest reduction in CH<sub>4</sub> flux rates (>200 kg CH<sub>4</sub>-C ha<sup>-1</sup> yr<sup>-1</sup>) occurred in Hainan, Sichuan, Hubei, and Guangdong provinces (Figure 3b), which are dominated by double-cropping rice systems with warm weather and high-clay soils; the lowest reduction in CH<sub>4</sub> flux rates (<25 kg CH<sub>4</sub>-C ha<sup>-1</sup> yr<sup>-1</sup>) occurred in Tianjin, Hebei, Ningxia, Liaoning, and Gansu provinces (Figure 3b), which are dominated by single cropping systems with relatively cool weather and low-clay soils.

[21] Shifting water management from CF to MSD increased N<sub>2</sub>O emissions from almost all rice paddies in China, although the MSD-induced N<sub>2</sub>O flux rates varied from province to province (Table 2). The total N<sub>2</sub>O emissions from the Chinese rice paddies under CF and MSD conditions were 290–410 and 420–610 Gg N<sub>2</sub>O-N yr<sup>-1</sup>, respectively (Table 2). Taking mean values, changing water management to MSD increased the rice paddy N<sub>2</sub>O flux for China from 160 Gg N<sub>2</sub>O-N yr<sup>-1</sup>, equivalent to 2–3% of global total anthropogenic N<sub>2</sub>O emissions [*Prather et al.*, 2001]. The MSD-induced increase in N<sub>2</sub>O fluxes were high (>8.0 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>) in Jilin, Liaoning, Heilongjiang, and Xinjiang where the paddy soils contained relatively high organic matter content, and low (<3.0 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>) in Beijing, Tianjin, Hebei, Henan, Yunnan, Gansu,

and Ningxia (Figure 3c) where the soil organic matter (SOM) contents were relatively low.

[22] The aggregate radiative forcing of changes in CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O emissions due to changing water management was quantified as a CO<sub>2</sub>-equivalent flux (Table 3, Figure 4). With a 20-year time horizon, the nationally aggregated average flux was a net uptake of 9900 kg CO<sub>2</sub>-eq ha<sup>-1</sup> yr<sup>-1</sup> (equivalent to 81 Gg C yr<sup>-1</sup> uptake for all rice paddies). As the time horizon for the GWP analysis lengthened, the relative importance of reduced methane fluxes diminished, and the average flux dropped to a net uptake of 2000 kg CO<sub>2</sub>-eq ha<sup>-1</sup> yr<sup>-1</sup> for a 100-year horizon (~16 Gg C yr<sup>-1</sup> uptake for all rice paddies), and approximately zero net emission for a 500-year time horizon (Table 3).

#### 4. Discussion and Conclusions


[23] China's total rice paddy methane emissions with continuous flooding (6400–12,000 Gg CH<sub>4</sub>-C yr<sup>-1</sup>) were about 10–20% of recent estimates of global methane emissions from rice paddies [*Prather et al.*, 2001]; China's paddy area is about 20% of the global total paddy area [*Maclean et al.*, 2002]. Total rice paddy nitrous oxide

emissions for mid-season draining management ( $420\text{--}610\text{ Gg N}_2\text{O-N yr}^{-1}$ ) were about 10% of global  $\text{N}_2\text{O}$  emissions from cropland [Prather *et al.*, 2001]; China's paddy area is about 2.2% of total global cropland area [FAO, 2004]. Multiple cropping (including paddy and upland crop rotations) and heavy fertilizer use (up to  $240\text{ kg N ha}^{-1}\text{ yr}^{-1}$ ) probably account for proportionally high  $\text{N}_2\text{O}$  emissions. Total paddy  $\text{CO}_2$  emissions ranged from  $-3500$  to  $+12,000\text{ Gg C yr}^{-1}$ , with most losses

occurring in northern provinces, which have been more recently cultivated and have higher SOC values. Net  $\text{CO}_2$  emissions in eastern China (Fujian, Jiangxi, Anhui, Jiangsu, and Zhejiang provinces) were near zero. Cai [1996] quantified the effects of land use on SOC in these eastern provinces, based on mean SOC values reported in the Second Soil Survey of China. He found that mean paddy field SOC was about half that of natural vegetation soils, and mean upland crop field SOC was about half that of paddy fields. He estimated total SOC loss for the cropland soils ( $0\text{--}0.62\text{ m}$ ) in the region at  $850,000\text{ Gg C}$ , based on the assumption that soils supporting natural vegetation at the time of the Second Soil Survey were representative of all pre-agricultural soils. Losses of SOC typically occur during the first few decades of cultivation [e.g., Smith *et al.*, 1997; Li *et al.*, 1994], and most paddies in eastern China are old enough that their SOC may have stabilized at low levels.

[24] Shifting water management from continuous flooding to midseason drainage reduced total net  $\text{CO}_2$  emissions by  $0$  to  $-1500\text{ Gg CO}_2\text{-C yr}^{-1}$ , reduced total  $\text{CH}_4$  emissions from paddy rice fields in China by  $4200$  to  $4700\text{ Gg CH}_4\text{-C yr}^{-1}$ , and increased total  $\text{N}_2\text{O}$  emissions from paddy rice fields in China by  $130$  to  $200\text{ Gg N}_2\text{O-N yr}^{-1}$ . The response to changing water management was not spatially uniform, which has significant implications for the design of greenhouse gas mitigation strategies. Although the change in  $\text{CO}_2$  flux was small from an aggregate greenhouse gas perspective (Table 3), enhanced  $\text{CO}_2$  losses were greater in northern China (Figure 3a), where SOC content was generally higher. The reduction in  $\text{CH}_4$  emission per unit area was also generally higher in northern China (Figure 3b), as was the increase in  $\text{N}_2\text{O}$  emissions (Figure 3c). Evaluated as  $\text{CO}_2$ -equivalent fluxes using 100-year GWP values, shifting water management from continuous flooding to midseason drainage made northern China a net source, and southern China a net sink (Figure 4b). For a 20-year time horizon, most provinces are effectively net sinks (Figure 4a), while for a 500-year time horizon most provinces are effectively net sources, except along the Yangtze River and in the south (Figure 4c).

[25] Implementing alternative water management in areas with warm weather, high clay content, and low organic matter content caused the largest reduction in radiative forcing, particularly for the short time horizons where the impact of changes in methane emissions is more pro-



**Figure 4.** Net provincial  $\text{CO}_2$ -equivalent emissions for conversion from continuous flooding to mid-season draining, calculated using (a) 20-year, (b) 100-year, and (c) 500-year global warming potential conversion factors for  $\text{CH}_4$  and  $\text{N}_2\text{O}$  (see equation (1)) [Ramaswamy *et al.*, 2001]. Sign convention is a positive flux difference (red shades) represents an increase in flux to the atmosphere, and a negative flux difference (blue shades) represents a decrease in flux to the atmosphere. Tibet (Xizang) and Qinghai Provinces have very small paddy areas, and are reported here as no data; Taiwan was excluded from the analysis because of a lack of crop rotation data. See color version of this figure at back of this issue.



nounced (Figure 4, Table 3). Sichuan, Hubei, Hunan, Guangdong, Guangxi, Anhui, and Jiangsu provinces, which possess almost 60% of China's rice paddies and produce just over 60% of the rice, fall into this category. In northeastern China, Heilongjiang, Jilin, and Liaoning provinces produce ~10% of the annual rice harvest on ~10% of the paddy area. Switching to mid-season draining in this region led to net increase in greenhouse gas emissions on all timescales except the 20-year time horizon for Heilongjiang Province, which had a nearly neutral 20-year GWP (Figure 4, Table 3).


[26] Although mid-season drainage is now widespread in China and Japan, alternative water management regimes are not common for many of the rice producing countries in South and Southeast Asia, where they are still being evaluated by agronomists [Barker and Molle, 2004]. Although adopting changes in water management probably will be driven primarily by grain yield and water use, as was the case for China, quantifying impacts of alternative management practices on greenhouse gas emissions may play a role in decision making. These changes could include both switching from continuous flooding to mid-season draining and switching from rainfed to irrigated rice. The effects of management alternatives for mitigation are likely to vary when they are applied across climatic zones, soil types, or farming systems. Process-based models integrated with GIS databases can play an important role in linking management changes to biogeochemical cycles based on spatially differentiated information, and either target mitigation efforts to the most beneficial regions or evaluate spatial variability of greenhouse gas impacts of management changes.

[27] **Acknowledgment.** The research has been supported by NASA's Terrestrial Ecology Program (NAG5-12838 and NAG5-7631) and the NASA EOS-IDS program (NAG5-10135).

## References


- Barker, R., and F. Molle (2004), Irrigation management in rice-based cropping systems: Issues and challenges in Southeast Asia, *Ext. Bull.* 543, Food and Fertil. Technol. Cent., Taipei, Taiwan. (Available at [www.ffc.agnet.org](http://www.ffc.agnet.org)).
- Beijing Agricultural University (1992), *Cropland Management* (in Chinese), 286 pp., Agric. Press, Beijing.
- Cai, Z. C. (1996), Effect of land use on organic carbon storage in soils in eastern China, *Water Air Soil Pollut.*, 91, 383–393.
- Cai, Z. C., G. X. Xing, G. Y. Shen, H. Xu, X. Y. Yan, and H. Tsuruta (1999), Measurements of CH<sub>4</sub> and N<sub>2</sub>O emissions from rice paddies in Fengqiu, China, *Soil Sci. Plant Nutr.*, 45, 1–13.
- Cai, Z., S. Sawamoto, C. Li, G. Kang, J. Boonjawat, A. Mosier, and R. Wassmann (2003), Field validation of the DNDC model for greenhouse gas emissions in East Asian cropping systems, *Global Biogeochem. Cycles*, 17(4), 1107, doi:10.1029/2003GB002046.
- Central Radio and Television School of Agriculture (CRTSA) (1995), *Textbooks for Primary Agricultural and Technical Training* (in Chinese), 6 vols., China Agric. Publ. House, Beijing.
- Chen, G. X., G. H. Huang, B. Huang, J. Wu, K. W. Yu, H. Xiu, X. H. Xue, and Z. P. Wang (1995), CH<sub>4</sub> and N<sub>2</sub>O emission from a rice field and effect of Azolla and fertilization on them (in Chinese), *Chin. J. Appl. Ecol.*, 6, 378–382.
- Cole, V., C. Cerri, K. Minami, A. Mosier, N. Rosenberg, and D. Sauerbeck (1996), Agricultural options for mitigation of greenhouse gas emissions, in *Climate Change 1995: Impacts, Adaptations and Mitigation of Climate Change: Scientific-Technical Analyses*, edited by R. T. Watson, M. C. Zinyowera, and R. H. Moss, pp. 745–771, Cambridge Univ. Press, New York.
- Cui, D., H. Liu, J. Min, and J. He (1994), *Atlas of Phenology for Major Crops in China* (in Chinese), 177 pp., Meteorol. Press, Beijing.
- Food and Agricultural Organization (FAO) (2004), FAOSTAT agricultural database; <http://apps.fao.org/>, Rome.
- Frolking, S., J. Qiu, S. Boles, X. Xiao, J. Liu, Y. Zhuang, C. Li, and X. Qin (2002), Combining remote sensing and ground census data to develop new maps of the distribution of rice agriculture in China, *Global Biogeochem. Cycles*, 16(4), 1091, doi:10.1029/2001GB001425.
- Gao, L. Z., Z. Q. Jin, Y. Huang, H. Chen, and B. B. Li (1992), *Rice Cultivational Simulation-Optimization-Decision Making System: RCSODS* (in Chinese), Agric. Sci. and Tech. Publ. House, Beijing.
- Huang, G., Z. Zhang, and Q. Zhao (1997), *Crop Practice in Southern China* (in Chinese), China Agric. Publ. House, Beijing.
- Huke, R. E., and E. H. Huke (1997), *Rice Area by Type of Culture: South, Southeast, and East Asia*, 59 pp., Intl. Rice Res. Inst. (IRRI), Los Baños, Philippines.
- Institute of Soil Science (1986), *The Soil Atlas of China*, Acad. Sin., Cartogr. Publ. House, Beijing.
- Intergovernmental Panel on Climate Change (IPCC) (1997), *Guidelines for National Greenhouse Gas Inventories*, Org. for Econ. Coop. Dev., Paris.
- International Rice Research Institute (IRRI) (2004), RiceStat database, <http://www.irri.org/science/ricestat/index.asp>, Los Baños, Philippines.
- King, J. Y., W. S. Reebergh, K. K. Thieler, G. W. Kling, W. M. Loya, L. C. Johnson, and K. J. Nadelhoffer (2002), Pulse-labeling studies of carbon cycling in Arctic ecosystems: The contribution of photosynthates to methane emission, *Global Biogeochem. Cycles*, 16(4), 1062, doi:10.1029/2001GB001456.
- Li, C. (1995), Impact of agricultural practices on soil C storage and N<sub>2</sub>O emissions in 6 states in the U.S., in *Soil Management and Greenhouse Effect*, edited by R. Lal et al., pp. 101–112, Lewis Publ., Boca Raton, Fla.
- Li, C. (2000), Modeling trace gas emissions from agricultural ecosystems, *Nutr. Cycl. Agroecosyst.*, 58, 259–276.
- Li, C. (2001), Biogeochemical concepts and methodologies: Development of the DNDC model (in Chinese with English abstract), *Quat. Sci.*, 21, 89–99.
- Li, C., S. Frolking, and T. A. Frolking (1992), A model of nitrous oxide evolution from soil driven by rainfall events: 1. Model structure and sensitivity, *J. Geophys. Res.*, 97, 9759–9776.
- Li, C., S. Frolking, and R. C. Harriss (1994), Modeling carbon biogeochemistry in agricultural soils, *Global Biogeochem. Cycles*, 8, 237–254.
- Li, C., V. Narayanan, and R. Harriss (1996), Model estimates of nitrous oxide emissions from agricultural lands in the United States, *Global Biogeochem. Cycles*, 10, 297–306.
- Li, C., J. Qiu, S. Frolking, X. Xiao, W. Salas, B. Moore III, S. Boles, Y. Huang, and R. Sass (2002), Reduced methane emissions from large-scale changes in water management in China's rice paddies during 1980–2000, *Geophys. Res. Lett.*, 29(20), 1972, doi:10.1029/2002GL015370.
- Li, C., A. Mosier, R. Wassmann, Z. Cai, X. Zheng, Y. Huang, J. Tsuruta, J. Boonjawat, and R. Lantin (2004), Modeling greenhouse gas emissions from rice-based production systems: Sensitivity and upscaling, *Global Biogeochem. Cycles*, 18(1), GB1043, doi:10.1029/2003GB002045.
- Li, C., S. Frolking, and K. Butterbach-Bahl (2005), Carbon sequestration in arable soil is likely to increase nitrous oxide emissions, *Climatic Change*, in press.
- Li, Q. (1992), Water management of paddy soils, in *Paddy Soils of China* (in Chinese), edited by Q. Li, Sci. Press, Beijing.
- Liu, Z., and Z. Mu (1993), *Cultivation Systems in China* (in Chinese), Agric. Press, Beijing, China.
- Lu, Y. H., A. Watanabe, and M. Kimura (2002), Contribution of plant-derived carbon to soil microbial dynamics in a paddy rice microcosm, *Biol. Fertil. Soils*, 36, 136–142.
- Maclean, J. L., D. C. Dawe, B. Hardy, and G. P. Hettel (Eds) (2002), *Rice Almanac*, 3rd ed., 253 pp., CAB Intl., Oxon UK.
- Marland, G., and T. Boden (2004), Carbon Dioxide Information Analysis Center database, <http://cdiac.esd.ornl.gov/ftp/trends/emissions/prc.dat>, Oak Ridge Natl. Lab., Oak Ridge, Tenn.
- National Soil Survey Office of China (1997), *Soils in China* (in Chinese), 6 vols., Agric. Publ. House, Beijing.
- Oenema, O., G. Velthof, and P. Kuikman (2001), Technical and policy aspects of strategies to decrease greenhouse gas emissions from agriculture, *Nutr. Cycl. Agroecosyst.*, 60, 301–315.
- Paul, E. A., and F. E. Clark (1989), *Soil Microbiology and Biochemistry*, second ed., 273 pp., Elsevier, New York.
- Prather, M., et al. (2001), Atmospheric chemistry and greenhouse gases, in *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by J. T. Houghton et al., pp. 239–287, Cambridge Univ. Press, New York.

- Ramaswamy, V., O. Boucher, J. Haigh, D. Hauglustaine, J. Haywood, G. Myhre, T. Nakajima, G. Y. Shi, and S. Solomon (2001), Radiative forcing of climate change, in *Climate Change 2001: The Scientific Basis—Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by J. T. Houghton et al., pp. 239–287, Cambridge Univ. Press, New York.
- Robertson, G. P., E. A. Paul, and R. R. Harwood (2000), Greenhouse gases in intensive agriculture: Contributions of individual gases to the radiative forcing of the atmosphere, *Science*, *289*, 1922–1925.
- Sass, R. L., F. M. Fisher, Y. B. Wang, F. T. Turner, and M. F. Jund (1992), Methane emission from rice fields: The effect of flood water management, *Global Biogeochem. Cycles*, *6*, 249–262.
- Shen, S. (1998), *Soil Fertility in China* (in Chinese), Chinese Agric. Press, Beijing.
- Shen, Z. R., X. L. Yang, and Y. S. Pei (1998), Enhancing researches on elevating efficiency of water use in Chinese agriculture, in *Strategies Against Water Crisis in Chinese Agriculture* (in Chinese), edited by Z. R. Shen and R. Q. Su, pp. 1–267, Chin. Agric. Sci. and Technol. Press, Beijing.
- Smith, P., et al. (1997), A comparison of the performance of nine soil organic matter models using datasets from seven long-term experiments, *Geoderma*, *81*, 153–225.
- Smith, P., K. W. Goulding, K. A. Smith, D. S. Powlson, J. U. Smith, P. Falloon, and K. Coleman (2001), Enhancing the carbon sink in European agricultural soils: Including trace gas fluxes in estimates of carbon mitigation potential, *Nutr. Cycl. Agroecosyst.*, *60*, 237–252.
- Stumm, W., and J. J. Morgan (1981), *Aquatic Chemistry: An Introduction Emphasizing Chemical Equilibria in Natural Waters*, 2nd ed., 780 pp., John Wiley, New York.
- Wassmann, R., H. U. Neue, R. S. Lantin, K. Makarim, N. Chareonsilp, L. V. Buendia, and H. Rennenberg (2000a), Characterization of methane emissions from rice fields in Asia: II. Differences among irrigation, rainfed, and deepwater rice, *Nutr. Cycl. Agroecosyst.*, *58*, 13–22.
- Wassmann, R., R. S. Lantin, H. U. Neue, L. V. Buendia, T. M. Corton, and Y. Lu (2000b), Characterization of methane emissions from rice fields in Asia: III. Mitigation options and future research needs, *Nutr. Cycl. Agroecosyst.*, *58*, 23–36.
- Watanabe, A., T. Takeda, and M. Kimura (1999), Evaluation of origins of CH<sub>4</sub> carbon emitted from rice paddies, *J. Geophys. Res.*, *104*, 23,623–23,629.
- Yagi, K., H. Tsuruta, K. Kanda, and K. Manami (1996), Effect of water management on methane emission from a Japanese rice field: Automated methane monitoring, *Global Biogeochem. Cycles*, *10*, 255–267.
- Yoshida, S. (1981), *Fundamentals of Rice Crop Science*, Intl. Rice Res. Inst., Los Baños, Philippines.
- Zheng, X. H., M. X. Wang, Y. S. Wang, R. X. Shen, X. J. Shangguan, J. Heyer, M. Kögge, H. Papen, J. S. Jin, and L. T. Li (1997), CH<sub>4</sub> and N<sub>2</sub>O emissions from rice paddies in southeast China, *Chin. J. Atmos. Sci.*, *21*, 167–174.
- Zheng, X., M. Wang, Y. Wang, R. Shen, J. Gou, J. Li, J. Jin, and L. Li (2000), Impacts of soil moisture on nitrous oxide emission from croplands: A case study on the rice-based agro-ecosystem in Southeast China, *Chemos. Global Change Sci.*, *2*, 207–224.
- 
- S. Boles, S. Frohling, C. Li, B. Moore III, and X. Xiao, Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, NH 03824, USA. (steve.frohling@unh.edu)
- Y. Huang, College of Resource and Environmental Sciences, Nanjing Agricultural University, Nanjing, China.
- J. Qiu, Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Science, Beijing, China.
- W. Salas, Applied Geosolutions, LLC, Durham, NH 03824, USA.
- R. Sass, Department of Ecology and Evolutionary Biology, Rice University, Houston, TX 77005, USA.




**Figure 1.** Comparison between observed and DNDC-modeled CH<sub>4</sub> and N<sub>2</sub>O fluxes from a paddy rice field applied with mid-season drainage in Wu County, Jiangsu Province, China, in 1995. DNDC captured the episodes of CH<sub>4</sub> emission depressions and N<sub>2</sub>O emission increases during the soil drying time periods by tracking the soil oxygen diffusion, CH<sub>4</sub> oxidation, labile organic matter decomposition, and stimulated nitrification and denitrification fueled by the increased ammonium and nitrate production due to the conversions of soil anaerobic to aerobic conditions driven by the mid-season drainage. (Field data were adopted from *Zheng et al.* [1997]).





**Figure 3.** Change, due to conversion from continuous flooding to mid-season draining, in annual paddy flux rates of (a) CO<sub>2</sub>, (b) CH<sub>4</sub>, and (c) N<sub>2</sub>O. Each provincial value is the difference between the mean for all counties of high and low values for mid-season draining and the mean for all counties of high and low values for continuous flooding. Sign convention: a positive flux difference (red shades) represents an increase in flux to the atmosphere, and a negative flux difference (blue shades) represents a decrease in flux to the atmosphere. For all provinces, the water management conversion reduced mean CH<sub>4</sub> flux and increase mean N<sub>2</sub>O flux. Tibet (Xizang) and Qinghai Provinces have very small paddy areas, and are reported here as no data; Taiwan was excluded from the analysis because of a lack of crop rotation data.



**Figure 4.** Net provincial CO<sub>2</sub>-equivalent emissions for conversion from continuous flooding to mid-season draining, calculated using (a) 20-year, (b) 100-year, and (c) 500-year global warming potential conversion factors for CH<sub>4</sub> and N<sub>2</sub>O (see equation (1)) [Ramaswamy *et al.*, 2001]. Sign convention is a positive flux difference (red shades) represents an increase in flux to the atmosphere, and a negative flux difference (blue shades) represents a decrease in flux to the atmosphere. Tibet (Xizang) and Qinghai Provinces have very small paddy areas, and are reported here as no data; Taiwan was excluded from the analysis because of a lack of crop rotation data.